MODEL PRACTICE TEST PAPER - II MATHEMATICS CLASS 12 - CBSE 2011

Time: 3 hrs Max. Marks: 100

General Instructions:

- All questions are compulsory
- The question paper consists of 29 questions divided into three sections A,B and C. Section A contains 10 questions of 1 mark each, Section B contains 12 questions of 4 marks each and section C contains 07 questions of 6 marks each.

3.

Section - A (Questions 1 - 10 carry one mark each)

- 1. $\int \frac{\log(\sin x)}{\tan x} dx$ 2. Write the principal value of $\cos^{-1} \cos(\frac{7\pi}{6})$
- 3. If $|\vec{a}| = \sqrt{3}$, $|\vec{b}| = 2$ and $\vec{a} \cdot \vec{b} = 3$, find the angle between \vec{a} and \vec{b}
- 4. Write down the equation of a line parallel to the line $\frac{x-2}{-3} = \frac{y+3}{2} = \frac{z+5}{6}$ and passing through the point
- 5. If matrix A = (1 2 3), write AA', where A' is the transpose of A
- 6. Evaluate $\int Sin4xcos3x dx$
- 7. Write the order and degree of differential equation $\left(\frac{d^2y}{dx^2}\right)^2 + \left(\frac{dy}{dx}\right)^3 + 2y = 0$
- 8. Find the value of p if $(2\hat{\imath}+6\hat{\jmath}+27\hat{k}) \times (\hat{\imath}+3\hat{\jmath}+p\hat{k}) = \vec{0}$ 9. Evaluate : $\begin{bmatrix} 2\cos\theta & -2\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$
- 10. Form the differential equation of the family of curves $y = a \cos(x+b)$, where a and b are arbitrary constants.

Section - B (Questions 11 – 22 carry four marks each)

- Questions 11 22 carry rour marks each, $1 \quad 1+p \quad 1+p+q$ 11. Using the properties of determinants, prove the following: $[2 \quad 3+2p \quad 1+3p+2q]=1$ $3 \quad 6+3p \quad 1+6p+3q$
- 12. If $y = (\sin x)^x + \sin^{-1} \sqrt{x}$, find $\frac{dy}{dx}$
- 13. Form a differential equation of the family of circles touching the x-axis at origin.
- 14. Solve the following for x: $\tan^{-1} x + 2\cot^{-1} x = \frac{2\pi}{3}$ or, Prove that $2\tan^{-1}\frac{1}{2} + \tan^{-1}\frac{1}{7} = \tan^{-1}\frac{31}{17}$ 15. Find the value of α so that the lines $\frac{1-x}{3} = \frac{7y-14}{2\alpha} = \frac{5z-10}{11}$ and $\frac{7-7x}{3\alpha} = \frac{y-5}{1} = \frac{6-z}{5}$ are perpendicular to each other.
- 16. Find the equation of the tangent to the curve $x^2+3y=3$, which is parallel to the line y-4x+5=0

Find the intervals in which the function f given by $f(x) = \sin x + \cos x$, $0 \le x \le 2\pi$, is strictly increasing or strictly decreasing.

- 17. Using properties of definite integral, evaluate : $\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx$, or, Evaluate $\int \frac{dx}{\sqrt{(5 4x 2x^2)}}$
- 18. Solve the differential equation : $\frac{dy}{dx} + y = \cos x \sin x$ 19. If f: N \rightarrow N be defined by f(n) = $\frac{n+1}{2}$, if n is odd = $\frac{n}{2}$, if n is even

Find whether the function f is bijective.

- 20. The scalar product of the vector $\vec{i} + \vec{j} + \vec{k}$ with unit vector along sum of vectors $2\vec{i} + 4\vec{j} 5\vec{k}$ and $\mu \vec{i} + 2\vec{j} + 3\vec{k}$ is equal to one. Find the value of μ
- 21. Let * be the Binary operation on N given by a*b = LCM of a and b. Find the value of 20*16. Is * (i) Commutative (ii) Associative.
- 22. Prove that the relation R in the set $A = \{1,2,3,4,5\}$ given by $R = \{(a,b) : |a-b| \text{ is even}\}$, is an equivalence relation.

Section - C (Questions 23 – 29 carry Six marks each)

23. Find the point on the curve $y^2=2x$ which is at a minimum distance from the point (1,4)

24. Obtain inverse of the following matrix using elementary operations $\begin{bmatrix} 2 & 3 & 0 \\ 2 & 3 & 0 \end{bmatrix}$

25. Evaluate $\int_0^{\pi} \frac{x \, dx}{a^2 \cos^2 x + b^2 \sin^2 x}$ 26. Find the area of the region enclosed between the two circles $x^2 + y^2 = 9$ and $(x-3)^2 + y^2 = 9$

27. Evaluate $\int \frac{x^2}{x^4 + x^2 + 1} dx$ or, Evaluate $\int_0^{\frac{\pi}{2}} (2 \log \sin x - \log \sin 2x) dx$

28. Find the equation of the plane through the point (-1,3,2) and perpendicular to each of the planes x + 2y + 3z = 5 and 3x + 3y + 3z = 0

29. A diet is to contain at least 80 units of vitamin A and 100 units of minerals. Two foods X and Y are available. Food X costs Rs.4 per unit and Food Y costs Rs.6 per unit. One unit of Food X contains 3 units of vitamin A and 4 units of minerals. One unit of Food Y contains 6 units of vitamin A and 3 units of minerals. Formulate this as a Linear Programming Problem and find graphically the minimum cost for diet that consists of mixture of these two foods and also meets the minimal nutritional requirements.

V.C.Srikanthan Chartered Engineer, MIE Cell: 9831071644

Email: contactvcs@rediffmail.com