



UNIVERSAL EDUCATION CENTRE

JAYANT SHARMA (94145-37474)

www.uecj4u.hpage.co.in

Time 3 to 3 $\frac{1}{2}$ Hrs

Maths 10 (Term 1)

M.M. 90

General Instructions:

- All questions are compulsory.
- The question paper consists of 34 questions divided into four sections A, B, C and D. Section A comprises of 8 questions of 1 mark each, Section B comprises of 6 questions of 2 marks each. Section C comprises of 10 questions of 3 marks each and Section D comprises of 10 questions of 4 marks each.
- Question numbers 1 to 8 in Section A are multiple choice questions where you are to select on correct option out of the given four.
- There is no overall choice. However, internal choice has been provided in 1 question of two marks, 3 questions of three marks each and 2 questions of four marks each. You have to attempt only one of the alternatives in all such questions.
- Use of calculator is not permitted.
- An additional 15 minutes time has been allotted to read this question paper only.

SECTION A

[1] If the system of liner equations $x - ky = 2$ and $3x + 2y = -5$ has a unique solution, then the value of k is:-

- [a] $k = \frac{2}{3}$ [b] $k \neq -\frac{2}{3}$ [c] $k = \frac{3}{2}$ [d] $k \neq -\frac{3}{2}$

[2] If $\tan \theta = \frac{3}{4}$ then the value of $\frac{1 - \cos \theta}{1 + \cos \theta}$ is :-

- [a] $-\frac{1}{9}$ [b] $\frac{2}{9}$ [c] 1 [d] $\frac{1}{9}$

[3] If $\sin 3x = \cos [x - 26^\circ]$ and $3x$ is an actual angle, then the value of x is :-

- [a] 29° [b] 26° [c] 29° [d] 13°

[4] If $x = 2^3 \times 3 \times 5^2$, $y = 2^2 \times 3^3$, then HCF [x, y] is :-


- [a] 12 [b] 108 [c] 6 [d] 36

[5] If a positive integer p is divided by 3, then the remainder can be:-

- [a] 1 or 3 [b] 1, 2 or 3 [c] 0, 1 or 2 [d] 2 or 3

[6] If the given figure, the value of $\tan P - \cot R$ is:-

- [a] 1 [b] 0 [c] -1 [d] 2




[7] Construction of a cumulative frequency table is useful in determining the :-

- [a] Mean [b] Median [c] Mode [d] All the above


[8] In the given figure, if $\angle A = \angle D = 90^\circ$, $AD = 6$ cm, $CD = 8$ cm and $BC = 26$ cm then ar(ΔABC) is :-

- [a] 240 cm^2 [b] 48 cm^2 [c] 120^2 [d] 260 cm^2



SECTION – B

[9] Find the value of p and q in the given figure, if ABCD is a rectangle



[10] If α and $\frac{1}{\alpha}$ are the zeroes of the polynomial $p(x) = 4x^2 - 2x + k - 4$, then find the value of k

Or

Divide the polynomial $p(x) = 5 - 3x + 3x^2 - x^3$ by $g(x) = x^2 - x + 1$ and find the quotient and remainder

[11] Without actually performing the long division, state whether $\frac{39}{343}$ will have a terminating or non-terminating, repeating decimal expansion


[12] Find the value of k, if $\frac{\cos 35^\circ}{\sin 55^\circ} + \frac{2 \sin \theta}{\cos(90^\circ - \theta)} = \frac{k}{2}$



UEC (Raisinghnagar) Classes Online
Engaging Students Empowering Parents

Visit: www.uecj4u.hpage.co.in

[13] ABC is right angle triangle with $\angle ABC = 90^\circ$, $BD \perp AC$, $DM \perp BC$, and $DN \perp AB$. prove that $DM^2 = DN \times BC$.



[14] The following table gives production yield per hectare of wheat of 100 farms of village:-

Production (in kg/hect)	25-35	35-45	45-55	55-65	65-75	75-85
No. of farms	4	6	10	26	35	19

Write the above distribution to a more than type distribution.

SECTION - C

[15] Prove that $\frac{7\sqrt{7}}{4}$ is irrational.

Or

Prove that $(16 - 5\sqrt{7})$ is irrational.

[16] If one diagonal of a trapezium divides the other diagonal in the ratio 1:2. Prove that one of the parallel sides is double the other?

[17] Prove that: $\frac{\sin \theta}{\cot \theta + \operatorname{cosec} \theta} = 2 + \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta}$?

[18] The sum of the numerator and denominator, of a fraction is 8. If 3 is added to both the numerator and the denominator, the fraction become $\frac{3}{4}$. Find the fraction.

Or

Seven times a two digit number is equal to 4 times the number obtained by reversing the order of its digits. If the difference of digit is 3, find the number.

[19] If one zero of the polynomial $p(x) = 3x^2 - 8x + 2k + 1$ is seven times of other, then find the zeroes and the value of k.

[20] If $\sin \theta + \sin^2 \theta + \sin^3 \theta = 1$, prove that $\cos^6 \theta - 4\cos^4 \theta + 8\cos^2 \theta = 4$.



[21] Find the mean of the following data, using step-deviation method:-

Class Interval	0-20	20-40	40-60	60-80	80-100	100-120
Frequency	7	8	12	10	8	5

Or

Class Interval	0-20	20-40	40-60	60-80	80-100
Frequency	17	28	32	p	19

If the mean of the above data is 50, then find the value of p?

[22] Prove that $\tan\theta - \cot\theta = \frac{2\sin^2\theta - 1}{\sin\theta \cos\theta}$

[23] In ΔABC , if AD is the median, then show that $AB^2 + AC^2 = 2[AD^2 + BD^2]$.

[24] Find the median of the following data:

Class-interval	10-20	20-30	30-40	40-50	50-60	60-70	70-80
Frequency	12	30	34	65	46	25	18

SECTION-D

[25] Prove that, if a line is drawn parallel to one side of a triangle, to intersect the other two sides in distinct points the other two sides are divided into the same ratio.

Or

Prove that in a right triangle the square of the hypotenuse is equal to the sum of the square of the other two sides?

[26] If $x = a \sin\theta$, $y = b \tan\theta$. prove that $\frac{a^2}{x^2} - \frac{b^2}{y^2} = 1$.

[27] On dividing $3x^3 + 4x^2 + 5x - 13$ by a polynomial $g(x)$, the quotient and remainder are $3x + 10$ and $16x + 43$ respectively, Find the polynomial $g(x)$.

[

28] The fraction become $\frac{9}{11}$, if 2 is added to both the numerator and the denominator. If 3 is added to both the numerator and the denominator, it becomes $\frac{5}{6}$. Find the fraction.

[29] The following table shows the ages of the patients admitted in a hospital during a year:

AGE(in years)	5-15	15-25	25-35	35-45	45-55	55-65
No. of patients	6	11	21	23	14	5

Find the mode and mean of the data given above.

[30] The perpendicular from A on the side BC of the ΔABC intersects BC at D such that $DB = 3CD$ Prove that $2AB^2 = 2AC^2 + BC^2$.

[31] Draw the graph of following eqⁿ:-

$$2x + 3y = 12 \text{ and } x - y = 1$$

Shade the region between the two lines and x – axis. Also, determine the vertices of the triangle so formed.



[32] Prove that: $\frac{\cos\theta}{1-\tan\theta} + \frac{\sin^2\theta}{\sin\theta-\cos\theta} = \sin\theta + \cos\theta$
Or

Evaluate: $\frac{\sin^2\theta + \sin^2(90^\circ - \theta)}{3(\sec^2 61^\circ - \cot^2 29^\circ)} - \frac{3 \cot^2 30^\circ \sin^2 54^\circ \sec^2 36^\circ}{2(\operatorname{cosec}^2 65^\circ - \tan^2 25^\circ)}$.

[33] In a sports meet, the number of players in Football, Hockey and Athletics are 48,60,132, respectively. Find the minimum number of room required, if in each room the same number of player are to be seated and all of them being in the same sports ?

[34] The following distribution gives the daily income of 65 workers of a factory :-

Daily income (in Rs)	100-120	120 – 140	140 -160	160- 180	180- 200
No. of worker	14	16	10	16	9

Convert the distribution above to a more than type cumulative frequency distribution and draw its ogive.